

ORIGINAL ARTICLES

Effect of maternal education on the rate of childhood handicap

Sherine Shawky, Walled M. Milaat, Bahaa A. Abalkhalil, Nadia K. Soliman

ABSTRACT

Objectives: The objectives of this study were to determine the relation between maternal education and various maternal risk factors, identify the impact of maternal education on the risk of childhood handicap and estimate the proportion of childhood handicap that can be prevented by maternal education.

Methods: Data was collected from all married women attending the two major maternity and child hospitals in Jeddah during April 1999. Women with at least one living child were interviewed for sociodemographic factors and having at least one handicapped child. The risk of having a handicapped child and the population attributable risk percent were calculated.

Results: Some potential risk factors are dominant in our society as approximately 30% of women did not attend school and 84% did not work. Consanguineous marriages accounted for about 43%. Pre-marriage counseling was limited as only 10% of women counseled before marriage. The proportion of unemployment and consanguineous marriages decreased significantly by increase in maternal education level. Conversely, the proportion of women reporting pre-marriage counseling increased significantly by increase in maternal education level. Approximately, 7% of women reported having at least one handicapped child. The risk of having a handicapped child showed a significant sharp decline with increase in maternal education level. At least 25% of childhood handicap can be prevented by achieving female primary education and up to half of cases can be prevented if mothers finish their intermediate education.

Conclusions: Female education plays a major role in child health. The results of this study suggest investment in female education, which would have substantial positive effects in reducing incidence of childhood handicap in Jeddah.

Saudi Medical Journal 2001; Vol. 22 (1): 39-43

Childhood handicap is a major public health concern especially in developing countries as over three quarters of the world's handicapped children under 15 years of age are estimated to live in this area.¹ It is very difficult to define where child health begins. In dealing with child health, one should recognize that it is an outcome of many intervening factors that start as early as the parent's early childhood and adolescence. In the last decades, this discipline has developed tremendously. The health professionals have begun to move from a narrow concept of maternal health to a more inclusive one that recognizes the role of much more fundamental issues substantially influencing child health.² Among these conditions, are all factors affecting maternal development that are consequently reflected on her offspring. The economic, social, cultural and educational environment in which girls grow to womanhood, marry and child-bear affects their future reproductive health and children.

In the Arab World, some potential risk factors issue of the traditions and beliefs still exist. Women's illiteracy, early marriage, multiparity and consanguineous marriages are still dominant.³⁻⁶ Saudi Arabia has undergone tremendous economic development in the recent years. Consequently, there has been marked improvement in the health care services and frank decline in infant morbidity and mortality. Development of health care services has emerged the problem of handicapped children as more feeble children survive. The prevalence of major disabilities was estimated to be around 4%.⁷ Female education has advanced and more girls are urged to attend schools. However, the level of illiteracy among women remains elevated and about 39% of girls do not attend schools.⁸ Also traditionally, girls marry at a young age,^{4,6} which interrupts their education at an early stage. This phenomenon is not unique for the Kingdom but is dominant in the Middle East Area.⁴⁻⁶ Female education has been previously recognized as a major risk factor for child health. At present, our understanding of risk factors are mostly derived from studies in the Western Countries where maternal illiteracy does not exist. Few experiences come from the Third World Area and need to be strengthened through research.

The objectives of this study were to: 1) Determine the relation between maternal education and various maternal risk factors in Jeddah; 2) Estimate the impact of maternal education level on the risk of childhood handicap and 3) Calculate the population attributable risk percent (PAR%) of childhood handicap that can be eliminated by maternal education.

Methods. Study site and data source. Jeddah city is one of the largest cities of the Kingdom of Saudi Arabia with a population of 2.1 million.⁹ In Jeddah, there are two major governmental maternity and child hospitals that drain around 60%³ of women of low and medium social class in Jeddah City. These two hospitals contain 290 inpatient beds and serve around 2000 patients per month. A cross-sectional study was performed in these two maternity hospitals to collect data on all married women with at least one living child during April 1999 either as patient (inpatient or outpatient) or healthy (who accompanied their child, any family member or a friend). Two trained medical students, under direct supervision of the medical staff, using structured questionnaires, interviewed women who accepted participation in the study in each hospital. A pilot study was carried out to test the questionnaire. Information on sociodemographic factors and having a handicapped child were collected. A total of 914 women were identified. In all, 826 (90%) were eligible for the study a

From the Department of Community Medicine and Primary Health Care, Faculty of Medicine and Allied Health Sciences, King AbdulAziz University, Jeddah. Received 1st July 2000. Accepted for publication in final form 12th September 2000 Address correspondence and reprint request to: Dr. Sherine Shawky, PO Box 115, Jeddah 21411, Kingdom of Saudi Arabia. Tel. +966 (2) 6318318 Fax. +966 (2) 632 3142. E-mail: shshawky@yahoo.com

References

1. Helander EAS. Prejudice and dignity: An introduction of community-based rehabilitation, New York: USA; United Nations Development Program 1993.
2. Sai FT, Nassim J. The need for reproductive health approach. Int J Gynecol Obstet 1989; suppl. 3: 103-113.

3. Milaat WA, Du V, Florey CH. Perinatal mortality in Jeddah, Saudi Arabia. *Int J Epid* 1992; 21: 82-90.
4. Harfouche JK, Verhostrate LJ. Child Health. The state of child health in the Eastern Mediterranean region. WHO/EMRO Technical Publication. Alexandria: Egypt; 1983; 9: 131-163.
5. El Hamamsy L. Early marriage & reproduction in two Egyptian villages. Occasional Monograph, The Population Council/UNFPA. Cairo: Egypt; 1994.
6. Shawky S, Milaat WA. Early Teenage Marriage and Subsequent Pregnancy Outcomes. Accepted for publication.
7. Al-Turaiki MH. National project for disability rehabilitation. King Fahad National Library. Riyadh: Kingdom of Saudi Arabia; 1997.
8. United Nations Population Fund. The state of the world population 1997. Geneva: UNFPA; 1997.
9. Milaat W. Public Health around the World: Saudi Arabia. *Public Health Medicine* 1999; 1: 34-38.
10. Seagar J, Olson A. Women in the World: An International Atlas. New York: Simon & Schuster; 1986. p.24.
11. Bicego GT, Boerma JT. Maternal education and child survival: A comparative study of survey data from 17 countries. *Soc Sci Med* 1993; 36: 1207-1227.
12. Becerra J, Atrash H, Perez N, Saliceti JA. Low birthweight and infant mortality in Puerto Rico. *Am J Public Health* 1993; 83: 1572-1576.
13. Victora CG, Huttly SRA, Barros FC, Lombardi C, Vaughan JP. Maternal education in relation to early and late child health outcomes: Findings from a Brazilian cohort study. *Soc Sci Med* 1992; 889-905.
14. Adetunji JA. Infant mortality in Nigeria: Effects of place of birth, mother's education and region of residence. *J Biosoc SCI* 1994; 26: 469-477.
15. Kabagarama D, Mulford CL. The relation between women's education, nutrition, fertility, GNP per capita and infant mortality: Implications for the role of women in development. *Int Contemp Sociol* 1989; 26: 189-200.
16. Caldwell JC. Routes to low mortality in poor countries. *Popul Dev Rev* 1986; 12: 171-220.
17. Terra de Souza AC, Cufino E, Peterson KE, Gardner J, Vasconcelos do Amaral MI and Ascherio A. Variations in infant mortality rates among municipalities in state of Ceara, Northeast Brazil: An ecological analysis. *IJE* 1999; 28: 267-275.
18. Moerman ML. Growth of the birth canal in adolescent girls. *Am J Obstet and Gynecol*

1982; 143: 523-528.

19.Hayes L, Crovitz E and Durham NC. Adolescent Pregnancy. Southern Medical Journal 1979; 72: 869-874.

20.Dott AB, Fort AT. Medical and social factors affecting early teenage pregnancy. Am J Obstet and Gynecol 1976; 125: 532-536.

21.Al-Salem M, Rawashdeh N. Consanguinity in North Jordan: Prevalence and Pattern. J Biosoc Science 1993; 25: 553-556.

22.Gevi D, Nathan R, Freudilich M. Consanguinity and congenital heart disease in rural Arab population in Northern Israel. Human Hered 1986; 36: 213-217.

23.Hamamy HA, Al-Bayat N, A-Qubaisy W. Consanguineous matings in the Iraq urban population and the effect of pregnancy outcome on infant mortality. Iraq Med J 1986; 34: 65-80.

24.Saha N, Al-Shaykh F. Inbreeding levels in Sudan. Biosoc Science 1998; 20: 333-336.